

PPAR gamma Rabbit pAb

产品信息

产品名称	产品编号	规格
PPAR gamma Rabbit pAb	31269ES50	50 μ L
	31269ES60	100 μ L

产品简介

Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis.

产品性质

别名	CIMT1; GLM1; NR1C3; Nuclear receptor subfamily 1 group C member 3; Peroxisome proliferator activated nuclear receptor gamma variant 1; Peroxisome proliferator activated receptor gamma 1; Peroxisome Proliferator Activated Receptor gamma; Peroxisome proliferator-activated receptor gamma; PPAR gamma; PPAR-gamma; PPARG; PPARG_HUMAN; PPARG1; PPARG2; PPARgamma.
SwissProt ID	P37231
基因 ID	5468
免疫原	The antiserum was produced against synthesized peptide derived from human PPAR-gamma. AA range:78-127.
缓冲液	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
分子量	Calculated MW: 58 kDa; Observed MW: 58 kDa
克隆类型	Polyclonal Antibody
宿主	Rabbit
物种反应	Human, Mouse, Rat
推荐应用	WB, IHC, ICC/IF, ELISA
纯化方式	Affinity Purified

应用

稀释比 WB: 1/500-1/1000; IHC: 1/50-1/100; IF: 1/50-1/200; ELISA: 1/10000

储存条件


-25~-15°C保存

注意事项

1. 避免反复冻融。
2. 本产品仅作科研用途。
2. 为了您的安全和健康，请穿实验服并佩戴一次性手套操作。


产品数据

Western blot



Western blot analysis of PPAR gamma in HuvEc lysates using PPAR gamma antibody.

Immunohistochemistry



Immunohistochemistry analysis of paraffin-embedded rat spleen tissue using PPAR γ antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Sample with blocking peptide on the right.